

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 85301895.0

(51) Int. Cl.⁴: **C 07 D 417/12**
A 61 K 31/44

(22) Date of filing: 19.03.85

(30) Priority: 21.03.84 PC T/JP84/00117
14.09.84 PC T/JP84/00445

(43) Date of publication of application:
25.09.85 Bulletin 85/39

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: Takeda Chemical Industries, Ltd.
27, Doshomachi 2-chome Higashi-ku
Osaka-shi Osaka, 541(JP)


(72) Inventor: Meguro, Kanji
24-311, 1 Kotobuki-cho
Nishinomiya-shi Hyogo 662(JP)

(72) Inventor: Fujita, Takeshi
13-15, Nagaodai 1-chome
Takarazuka-shi Hyogo 665(JP)

(74) Representative: Laredo, Jack Joseph et al,
Elkington and Fife High Holborn House 52/54 High
Holborn
London, WC1V 6SH(GB)

(54) Thiazolidinedione derivatives, their production and use.

(57) Thiazolidinedione derivatives of the general formula:



[wherein R¹ and R² are the same or different and each represent hydrogen or a lower alkyl group; R³ is hydrogen or an acyl group; n is 0 or 1] and salts thereof are novel compounds, which exhibit in mammals blood sugar- and lipid-lowering activity, and are of value as a therapeutic agent for diabetes and therapeutic agent for hyperlipemia.

- 1 -


Thiazolidinedione Derivatives, Their Production and Use

The present invention relates to novel thiazolidinedione derivatives having blood sugar- and lipid-lowering activity, to a process for producing the same, and to pharmaceutical compositions containing the same.

5 A great variety of biguanide and sulfonylurea compounds have been conventionally employed as a therapeutic agent for diabetes. However, biguanide compounds cause lactic acidosis and therefore, are hardly used at present, while sulfonylurea compounds exhibit potent blood sugar lowering activity but
10 often bring about serious hypoglycemia, thus requiring precautions for use. The present inventors, after extensive research to find a compound having blood sugar lowering activity which is free from such drawbacks, found out novel thiazolidinedione derivatives having excellent blood sugar-
15 and lipid-lowering activity.

The present invention relates to:

1. Thiazolidinedione derivatives of the general formula:



[wherein R¹ and R² are the same or different and each represent hydrogen or a lower alkyl group; R³ is hydrogen or an acyl
25 group; n is 0 or 1],

2. A process for producing thiazolidinedione derivatives of the general formula (I), characterized in that said process

The thiazolidinedione derivatives of the general formula (I) are amphoteric compounds having acid nitrogen in the thiazolidine ring and alkaline nitrogen in the pyridine ring, and exist both in acid and base salts. Referring to salts of the thiazolidinedione derivatives (I), examples of the acid salts include inorganic acid salts, such as hydrochlorides, hydrobromides, sulfates, phosphates and methanesulfonates, and organic acid salts, such as toluenesulfonates, oxalates, malonates, maleates, fumarates, succinates, tartrates and malates, while examples of the base salts include metal salts, such as sodium salts, potassium salts, aluminum salts, magnesium salts and calcium salts.

As specific examples of the thiazolidinedione derivatives of the general formula (I) and their salts, there may be mentioned the following compounds:

- 5-{4-[2-Hydroxy-2-(5-methyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-Acetoxy-2-(5-methyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[3-Hydroxy-2-(3-methyl-2-pyridyl)propoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[3-Acetoxy-2-(3-methyl-2-pyridyl)propoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[3-Benzoyloxy-2-(3-methyl-2-pyridyl)propoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-(5-Ethyl-2-pyridyl)-2-hydroxyethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-(5-Ethyl-2-pyridyl)-2-formyloxyethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-Acetoxy-2-(5-ethyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-(5-Ethyl-2-pyridyl)-2-propionyloxyethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-(5-Ethyl-2-pyridyl)-2-butyryloxyethoxy]benzyl}-2,4-thiazolidinedione,
- 5-{4-[2-(5-Ethyl-2-pyridyl)-2-isobutyryloxyethoxy]benzyl}-2,4-thiazolidinedione,

- 5-{4-[2-(5-Ethyl-2-pyridyl)-2-valeryloxyethoxy]benzyl}-
2,4-thiazolidinedione,
5-{4-[2-Benzoyloxy-2-(5-ethyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione,
5 5-{4-[2-(5-Ethyl-2-pyridyl)-3-hydroxypropoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[3-Acetoxy-2-(5-ethyl-2-pyridyl)propoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[2-Hydroxy-2-(2-pyridyl)ethoxy]benzyl}-2,4-
10 thiazolidinedione,
5-{4-[2-Acetoxy-2-(2-pyridyl)ethoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[2-Propionyloxy-2-(2-pyridyl)ethoxy]benzyl}-2,4-
thiazolidinedione,
15 5-{4-[2-Benzoyloxy-2-(2-pyridyl)ethoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[3-Hydroxy-2-(2-pyridyl)propoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[3-Acetoxy-2-(2-pyridyl)propoxy]benzyl}-2,4-
20 thiazolidinedione,
5-{4-[2-Hydroxy-2-(5-methyl-2-pyridyl)ethoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[2-Acetoxy-2-(5-methyl-2-pyridyl)ethoxy]benzyl}-2,4-
thiazolidinedione,
25 5-{4-[2-(5-Methyl-2-pyridyl)-2-propionyloxyethoxy]benzyl}-
2,4-thiazolidinedione,
5-{4-[2-Benzoyloxy-2-(5-methyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione,
5-{4-[3-Hydroxy-2-(5-methyl-2-pyridyl)propoxy]benzyl}-
30 2,4-thiazolidinedione,
5-{4-[2-Hydroxy-2-(3-methyl-2-pyridyl)ethoxy]benzyl}-2,4-
thiazolidinedione,
5-{4-[2-Acetoxy-2-(3-methyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione,
35 5-{4-[2-Benzoyloxy-2-(3-methyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione,

0155845

5-{4-[2-Hydroxy-2-(4-methyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione,

5-{4-[2-Acetoxy-2-(4-methyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione,

5 5-{4-[2-Hydroxy-2-(4,6-dimethyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione, and

5-{4-[2-Acetoxy-2-(4,6-dimethyl-2-pyridyl)ethoxy]benzyl}-
2,4-thiazolidinedione.

The reaction of a compound of the general formula (II)
10 with thiourea is normally conducted in a solvent, such as
alcohols (e.g., methanol, ethanol, propanol, 2-propanol,
butanol, isobutanol, 2-methoxyethanol, etc.), dimethylsulfoxide
and sulfolane. The reaction temperature normally ranges
15 from 20°C to 180°C, preferably from 60°C to 150°C. The
amount of thiourea to be used is 1 to 2 moles per mole of
the compound (II). In this reaction, hydrogen halide is
formed as a by-product as the reaction proceeds, and the
reaction may be carried out in the presence of a deacidifying
20 agent added, such as sodium acetate and potassium acetate,
in order to capture such hydrogen halide. The deacidifying
agent is normally used at a ratio of 1 to 1.5 moles per mole
of the compound (II). By means of such reactions, the compound
(III) is produced, and can be isolated, if desired, or may
25 be subjected to the next hydrolysis step directly without
isolation of the compound (III).

Hydrolysis of the compound (III) is normally carried
out in an appropriate solvent and in the presence of water
and a mineral acid. As the solvent, there may be mentioned
such solvents as may be used in the above-described reaction
30 of the compound (II) with thiourea. Examples of the mineral
acid include hydrochloric acid, hydrobromic acid, sulfuric
acid and so forth, and its amount to be used is 0.1 to 10
moles per mole of the compound (III), preferably 0.2 to 3
moles. The addition amount of water is normally a large
35 excess amount against 1 mole of the compound (II). This
reaction is normally conducted under warming or heating,

and the reaction temperature is normally 60 to 150°C. The reaction time ranges normally from several hours to ten-odd hours.

5 By means of this reaction, there can be obtained the compound of the general formula (I) where R^3 is hydrogen (hereinafter referred to in some instances as "hydroxyl compound (I')").

The above-mentioned hydroxyl compound (I') may be subjected to the following acylation reaction, if necessary.

10 The acylation reaction of the hydroxyl compound (I') is normally carried out by the action of an acylating agent in an appropriate solvent and in the presence of a base. Examples of the said solvent include esters, such as ethyl acetate, aromatic hydrocarbons, such as benzene, toluene and
15 xylene, ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran and dioxane, ketones, such as acetone and methyl ethyl ketone, and chlorinated hydrocarbons, such as dichloromethane, chloroform and carbon tetrachloride, as well as dimethylformamide, etc. As the acylating agent, there may
20 be mentioned formic acid, acid anhydrides or acid halides of aliphatic, aromatic-aliphatic or aromatic carboxylic acids, and the like. Examples of the said aliphatic carboxylic acids include those having 2 to 6 carbon atoms, such as acetic acid, propionic acid, butyric acid, isobutyric acid, valeric acid,
25 isovaleric acid and hexanoic acid, and as examples of the said aromatic-aliphatic carboxylic acids, there may be mentioned those having 8 to 9 carbon atoms, such as phenylacetic acid and phenylpropionic acid, while examples of the said aromatic carboxylic acids include those having 7 to 8 carbon atoms,
30 such as benzoic acid and p-methylbenzoic acid, whereby for example halogen (e.g., fluorine, chlorine, bromine, etc.), alkoxy (e.g., methoxy, ethoxy, etc.) and trifluoromethyl groups may further be substituted for hydrogen atoms on these aromatic nuclei.

35 The amount of the acylating agent to be used is normally 1 to 10 moles per mole of the hydroxyl compound (I'), preferably

1 to 2 moles. Examples of the base include amines, such as pyridine and triethylamine, and carbonates and hydrogen-carbonates, such as sodium carbonate, potassium carbonate, sodium hydrogencarbonate and potassium hydrogencarbonate. The base is normally used in an equimolar amount or a large excess amount against the acylating agent. In cases in which pyridine is used as a base, a large excess amount of pyridine employed can also serve as a solvent. This reaction is normally carried out at a temperature of -20°C to 40°C, and the reaction time normally ranges from 10 minutes to 24 hours. By means of this reaction, there can be obtained the compound of the general formula (I) where R³ is an acyl group [hereinafter referred to in some instances as "acyl compound (I)"].

The thiazolidinedione derivatives (I), if necessary, can be converted into salts by reacting with acid or base in accordance with a conventional method.


The thiazolidinedione derivatives and their salts as obtained in the above manner can be isolated and purified by known separation/purification means, such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, phase transfer and chromatography.

The compounds (I) of the present invention and their salts, in mammals (e.g., mice, dogs, cats, monkeys and man), demonstrate excellent blood sugar- and lipid-lowering activity, and in addition, are less toxic, being low both in acute and subacute toxicities; for example, 5-{4-[2-hydroxy-2-(6-methyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione, 5-{4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]benzyl}-2,4-thiazolidinedione, 5-{4-[2-(5-ethyl-2-pyridyl)-2-hydroxy]ethoxybenzyl}-2,4-thiazolidinedione, 5-{4-[2-(5-ethyl-2-pyridyl)-3-hydroxypropoxy]benzyl}-2,4-thiazolidinedione, 5-{4-[2-acetoxy-2-(6-methyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione and 5-{4-[3-acetoxy-2-(5-ethyl-2-pyridyl)propoxy]benzyl}-2,4-thiazolidinedione were all found to exhibit an LD₅₀ value of not less than 5000 mg/kg in rats when administered

orally. These suggest that the thiazolidinedione derivatives (I) and their salts are useful for the treatment of hyperlipemia, diabetes and their complications. With reference to the method of administration, they are normally usable orally for example in the form of tablets, capsules, powders, granules, etc., but can also be administered parenterally in the form of injectable solutions, suppositories, pellets, etc., as the case may be. In the case of their being employed as a therapeutic agent for diabetes of humans, they can be normally administered orally in daily doses of 0.05 mg to 10 mg/kg or parenterally in daily doses of 0.01 mg to 10 mg/kg, and in the case of their being utilized as a therapeutic agent for hyperlipemia of humans, they can be normally given orally in daily doses of 0.05 mg to 10 mg/kg or parenterally in daily doses of 0.01 mg to 10 mg/kg, whereby it is desirable to administer such quantities once daily or intermittently twice to four times weekly.

The starting compound (II) of the present invention can be produced for example by the following methods.

1) Production of the compound of the general formula (II) where n is 0;



[wherein R¹, R² and X are as defined hereinbefore; R^{4'} is as defined by the acyl group represented by R³ or R^{4'}; R^{5'} is as


0155845

defined by the lower alkyl group represented by R^5

The oxidation reaction of the compound (IV) to the compound (V) can be carried out easily by reacting the compound (IV) with hydrogen peroxide or an organic peracid.

5 Examples of such an organic peracid include performic acid, peracetic acid, pertrifluoroacetic acid, perbenzoic acid, m-chloroperbenzoic acid, etc., and this oxidation reaction can be conducted in accordance with the general methods known per se. The acylation reaction of the compound (V) to the
10 compound (II') is carried out by reacting the compound (V) with an acylating agent, and is easily conducted normally by heating with an acid anhydride or acid halide at 80°C to 150°C. The hydrolysis reaction of the compound (II') to the compound (II'') can be conducted by an ordinary method using
15 sodium hydroxide or potassium hydroxide.

ii) Production of the compound of the general formula (II) where n is 1:



30 [wherein each of the symbols is as defined hereinbefore].

The reaction of the compound (VI) to the compound (VIII) is carried out by allowing the compounds (VI) and (VII) to undergo condensation in the presence of sodium hydride. This reaction can be conducted in a solvent such as dimethyl-
35 formamide and tetrahydrofuran at -10°C to 20°C. The subsequent reaction of the compound (VIII) to the compound (IX) is

carried out, for example, by subjecting the compound (VIII) to catalytic reduction in an ordinary manner with use of palladium carbon as a catalyst. The reaction of the compound (IX) to the compound (II'') is conducted by diazotizing the compound (IX) in the presence of a hydrogen halide (HX) and further subjecting the resulting diazotized compound to the so-called Meerwein arylation reaction which involves the reaction with acrylic acid or its esters (X) in the presence of a copper catalyst (e.g., cuprous oxide, cupric oxide, cuprous chloride, cupric chloride, cuprous bromide, cupric bromide, etc.). The reaction of the compound (II'') to the compound (II''') is conducted by acylating the compound (II''). This reaction can be carried out in the same manner as the procedure of acylating the above-described compound (V).

The examples, reference examples and experiment example are described below to illustrate the present invention more specifically, but are not intended to limit the present invention.

Example 1

A mixture of methyl 3-{4-[2-acetoxy-2-(6-methyl-2-pyridyl)ethoxy]phenyl}-2-bromopropionate (3.2 g), thiourea (558 mg), sodium acetate (599 mg) and ethanol (30 ml) was heated under reflux for 4 hours, and then 6N-hydrochloric acid (30 ml) was added to the mixture, followed by heating under reflux for 16 hours. After neutralization with aqueous sodium hydrogencarbonate solution, the reaction mixture was extracted with chloroform, and the chloroform layer was washed with water and dried (MgSO₄). After the solvent was distilled off, the residue was chromatographed on a silica-gel (100 g) column and elution was effected with benzene-acetone (10:1, V/V) to give 5-{4-[2-hydroxy-2-(6-methyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione in the form of crystals. Yield of 0.95 g. Recrystallization from ethyl acetate-hexane produced colorless prisms. m.p. 154-155°C.

0155845

Elemental analysis, for $C_{18}H_{18}N_2O_4S$

Calcd.: C, 60.32; H, 5.06; N, 7.82

Found : C, 60.53; H, 5.24; N, 7.75

Example 2

5 (1) A mixture of methyl 2-bromo-3-{4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]phenyl}propionate (4.7 g), thiourea (875 mg), sodium acetate (943 mg) and ethanol (50 ml) was heated under reflux for 3 hours. The mixture was diluted with
10 water, neutralized with aqueous sodium hydrogencarbonate solution and extracted with ethyl acetate. The ethyl acetate layer was washed with water, dried ($MgSO_4$) and freed of the solvent to give 5-{4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]-benzyl}-2-imino-4-thiazolidinone in the form of crystals. Yield of 25 g. Recrystallization from chloroform-methanol
15 produced colorless prisms. m.p. 213-214°C.

Elemental analysis, for $C_{19}H_{21}N_3O_3S$

Calcd.: C, 61.44; H, 5.70; N, 11.31

Found : C, 61.14; H, 5.62; N, 10.99

(2) 5-{4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]benzyl}-2-imino-4-thiazolidinone (2.3 g) was dissolved in a mixture
20 of 2N-hydrochloric acid and ethanol (20 ml), and the solution was heated under reflux for 4 hours. After neutralization with aqueous sodium hydrogencarbonate solution, the solution was extracted with chloroform, and the chloroform layer was
25 washed with water and dried ($MgSO_4$). The solvent was distilled off to give 5-{4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]-benzyl}-2,4-thiazolidinedione in the form of crystals. Yield of 1.9 g. Recrystallization from ethanol produced colorless prisms. m.p. 182-183°C.

30 Elemental analysis, for $C_{19}H_{20}N_2O_4S$

Calcd.: C, 61.27; H, 5.41; N, 7.52

Found : C, 61.57; H, 5.49; N, 7.74

Example 3

By following the same procedure as described in Example 1,
35 methyl 3-{4-[2-acetoxy-2-(5-ethyl-2-pyridyl)ethoxy]phenyl}-

2-bromopropionate (10.0 g), thiourea (2.0 g) and sodium acetate (2.2 g) were allowed to undergo reaction in ethanol and hydrolysis was performed to give 5-{4-[2-(5-ethyl-2-pyridyl)-2-hydroxyethoxy]benzyl}-2,4-thiazolidinedione in the form of crystals. Yield of 6.2 g. Recrystallization from ethyl acetate-hexane produced colorless prisms. m.p. 129-130°C. Elemental analysis, for $C_{19}H_{20}N_2O_4S$
 Calcd.: C, 61.27; H, 5.41; N, 7.52
 Found : C, 61.36; H, 5.71; N, 7.08

Example 4

A mixture of methyl 2-bromo-3-{4-[2-(5-ethyl-2-pyridyl)-3-hydroxypropoxy]phenyl}propionate (12.2 g), thiourea (2.2 g), sodium acetate (2.4 g) and ethanol (100 ml) was heated under reflux for 3 hours, and 3N-hydrochloric acid (100 ml) was added to the mixture, followed by heating under reflux for 12 hours. After concentration, the reaction mixture was neutralized with aqueous sodium hydrogencarbonate solution and extracted with ethyl acetate. The ethyl acetate layer was washed with water, dried ($MgSO_4$) and then freed of the solvent. The residue was chromatographed on a silica-gel (200 g) column, and elution was effected with benzene-acetone (5:1, V/V) to give 5-{4-[2-(5-ethyl-2-pyridyl)-3-hydroxypropoxy]benzyl}-2,4-thiazolidinedione in the form of an oily substance. Yield of 4.8 g (42.9 %). The substance was dissolved in methanol (15 ml), and sodium methylate (28 % methanol solution, 2.9 g) was added to the solution, followed by addition of ethyl ether (150 ml) to give the sodium salt in the form of crystals. Yield of 4.8 g (40.7 %). Recrystallization from ethanol-ethyl ether produced colorless prisms. m.p. 237-238°C.

Elemental analysis, for $C_{20}H_{21}N_2O_4S \cdot Na$
 Calcd.: C, 58.81; H, 5.18; N, 6.86
 Found : C, 59.06; H, 4.94; N, 6.97

Example 5

A mixed solution of 5-{4-[2-hydroxy-2-(6-methyl-2-


pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione (350 mg), pyridine (5 ml) and acetic anhydride (0.2 ml) was allowed to stand at room temperature for 2 days, and then poured in water, followed by extraction with ethyl acetate. The ethyl acetate layer was washed with water, dried (MgSO_4) and freed of the solvent. The residue was chromatographed on a silica gel (20 g) column, and elution was effected with benzene-acetone (15:1, V/V) to give 5-{4-[2-acetoxy-2-(6-methyl-2-pyridyl)ethoxy]benzyl}-2,4-thiazolidinedione in the form of an oily substance. Yield of 360 mg (89.8 %). IR (neat): 1740, 1690, 1230 cm^{-1} . NMR (CDCl_3) δ : 2.11(3H,s), 2.51(3H,s), 3.02(1H,dd,J=14 and 9Hz), 3.41(1H,dd,J=14 and 4Hz), 4.37(2H,d,J=5Hz), 4.44(1H,dd,J=9 and 4Hz), 6.13(1H,t,J=5Hz), 6.83(2H,d,J=9Hz), 7.09(2H,d,J=9Hz), 7.0-7.7(3H,m).

Example 6

A mixture of 5-{4-[2-(5-ethyl-2-pyridyl)-3-hydroxypropoxy]benzyl}-2,4-thiazolidinedione sodium salt (408 mg), pyridine (10 ml) and acetic anhydride (0.15 ml) was stirred at room temperature for 8 hours, and poured into water, followed by extraction with ethyl acetate. The ethyl acetate layer was washed with water, dried (MgSO_4) and freed of the solvent. The residue was chromatographed on a silica-gel (20 g) column and elution was effected with benzene-acetone (10:1, V/V) to give 5-{4-[3-acetoxy-2-(5-ethyl-2-pyridyl)propoxy]benzyl}-2,4-thiazolidinedione in the form of an oily substance. Yield of 350 mg (81.8 %). IR(Neat): 1740, 1690, 1230 cm^{-1} . NMR (CDCl_3) δ : 1.24(3H,t,J=7Hz), 2.00(3H,s), 2.64(2H,q,J=7Hz), 3.03(1H,dd,J=14 and 9Hz), 3.45(1H,dd,J=14 and 9Hz), 3.60(1H,m), 4.31(2H,d,J=5Hz), 4.47(1H,dd,J=9 and 4Hz), 4.55(2H,d,J=5Hz), 6.80(2H,d,J=9Hz), 7.10(2H,d,J=9Hz), 7.20(1H,d,J=8Hz), 7.48(1H,dd,J=8 and 2Hz), 8.42(1H,d,J=2Hz).

Examples 7-9

By following the same procedure as described in Example 1, there were obtained the following compounds.



10

Example No.	R	melting point (°C)	Recrystallization solvent	yield (%)
7	3-CH ₃	203-204	DMF-water	83.8
8	H	162-164	ethanol	46.0
9	5-CH ₃	135-137	ethyl acetate-hexane	59.8

15

Example 10

20

By following the same procedure as described in Example 4, sodium salt of 5-(4-[2-(4,6-dimethyl-2-pyridyl)-2-hydroxyethoxy]benzyl)-2,4-thiazolidinedione was obtained. Yield: 72.7%. Recrystallization from ethanol-ether produced colorless crystals. m.p. 228-229°C.

25

Elemental analysis, for C₁₉H₁₉N₂O₄S·Na

Calcd.: C, 57.86; H, 4.86; N, 7.10

Found : C, 57.38; H, 4.69; N, 6.96

Example 11

30

A mixed solution of 5-(4-[2-(5-ethyl-2-pyridyl)-2-hydroxyethoxy]benzyl)-2,4-thiazolidinedione (500 mg), pyridine (5 ml) and propionic anhydride (0.3 ml) was allowed to stand over night at room temperature, and then poured into water, followed by extraction with ethyl acetate. The ethyl acetate layer was washed with water, dried (MgSO₄) and freed of the solvent. The residue was chromatographed on

35

a silica gel (30 g) column, and elution was effected with benzene-acetone (15:1, V/V) to give 5-{4-[2-(5-ethyl-2-pyridyl)-2-propionyloxyethoxy]benzyl}-2,4-thiazolidinedione in the form of an oily substance. Yield of 515 mg (89.7%).

5 IR(Neat): 1745, 1700, 1170 cm^{-1}

NMR (CDCl_3) δ : 1.13(3H,t,J=7Hz), 1.23(3H,t,J=7Hz),
2.40(2H,q,J=7Hz), 2.31(2H,q,J=7Hz), 3.02(1H,dd,
J=14 and 10Hz), 3.4(1H,m), 4.38(2H,d,J=5Hz),
4.40(1H,dd,J=10 and 4Hz), 6.18(1H,t,J=5Hz),
10 6.81(2H,d,J=9Hz), 7.10(2H,d,J=9Hz), 7.3~7.6(2H,m),
8.43(1H,d,J=2Hz), 9.2(1H, broad).

Example 12

15 Example of production of tablets:

(1) 5-{4-[2-(5-ethyl-2-pyridyl)-2-hydroxyethoxy]-benzyl}-2,4-thiazolidinedione	100 g
(2) Lactose	50 g
(3) Corn starch	15 g
20 (4) Carboxymethylcellulose calcium	44 g
(5) Magnesium stearate	1 g

210 g for 1000 tablets

The total amount each of the ingredients (1), (2) and (3) are kneaded with 30 g of the ingredient (4) in the presence of water, and the mixture is vacuum-dried and
25 granulated. The resulting granulates are mixed with 14 g of the ingredient (4) and 1 g of the ingredient (5) and compressed into tablets by use of a tablet compressing machine to manufacture 1000 tablets of 8 mm in diameter each
30 containing 100 mg of the ingredient (1).

Reference Example 1

(1) m-Chloroperbenzoic acid (70% of content, 7.1 g) was added to a solution of methyl 2-bromo-3-{4-[2-(6-methyl-2-pyridyl)ethoxy]phenyl}propionate (10.0 g) in methanol (100 ml) and the mixture was heated under reflux for 30 minutes.

After aqueous sodium thiosulfate solution was added to decompose the excess oxidizing agent, water was added to the reaction mixture, followed by extraction with ethyl acetate. The ethyl acetate layer was washed successively with water,

2N-potassium hydroxide and water in the mentioned order, dried (MgSO_4) and freed of the solvent by distillation to give a crude oily substance of methyl 2-bromo-3-{4-[2-(6-methyl-2-pyridyl)ethoxy]phenyl}propionate N-oxide. Yield of 10.1 g. IR (Neat): $1740, 1250 \text{ cm}^{-1}$.

(2) A mixture of methyl 2-bromo-3-{4-[2-(6-methyl-2-pyridyl)ethoxy]phenyl}propionate N-oxide (10.1 g) as obtained under (1) and acetic anhydride (50 ml) was heated at 110°C for 1 hour and concentrated under reduced pressure, and the residue was chromatographed on a silica-gel (200 g) column.

Elution was effected with benzene-acetone (50:1, V/V) to give methyl 3-{4-[2-acetoxy-2-(6-methyl-2-pyridyl)ethoxy]phenyl}-2-bromopropionate in the form of an oily substance. Yield of 3.3 g. IR (neat): 1740 cm^{-1} (broad). NMR (CDCl_3) δ : 2.13(3H,s), 2.53(3H,s), 3.14(1H,dd,J=14 and 7Hz), 3.40(1H,

dd, J=14 and 7Hz), 3.70(3H,s), 4.2-4.5(3H,m), 6.15(1H,t, J=6Hz), 6.8-7.6(7H,m).

Reference Example 2

- (1) 60% Oil-borne sodium hydride (2.8 g) was added little by little to a solution of 2-(3-methyl-2-pyridyl)-1,3-propylene glycol (10.0 g) and p-fluoronitrobenzene (8.45 g) in dimethylformamide (100 ml) under ice-cooling with stirring. After stirring under ice-cooling for 1 hour, the reaction solution was poured into ice-cold water, and the aqueous mixture was extracted with ethyl acetate. The ethyl acetate layer was washed with water, dried (MgSO₄) and freed of the solvent by distillation, and the residue was chromatographed on a silica-gel (200 g) column. Elution was effected with cyclohexane-ethyl acetate (3:1, V/V) to give 4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]nitrobenzene in the form of crystals. Yield of 9.6 g. Recrystallization from ethyl acetate-hexane produced colorless prisms. m.p. 135-136°C. Elemental analysis, for C₁₅H₁₆N₂O₄
Calcd.: C, 62.49; H, 5.59; N, 9.72
Found : C, 62.60; H, 5.69; N, 9.93
- (2) 4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]nitrobenzene (9.0 g) as obtained under (1) was dissolved in methanol (100 ml), and catalytic reduction was conducted at ambient temperature and at atmospheric pressure with use of 10% palladium-carbon. The catalyst was filtered out, and the solvent was distilled off. Then, the residue was dissolved in a mixture of methanol (80 ml), acetone (20 ml) and 47% aqueous hydrobromic acid (21.5 g), and a solution of sodium nitrite (2.4 g) in water (5 ml) was added dropwise to the solution at a temperature of not higher than 5°C under ice-cooling, followed by stirring at 5°C for 30 minutes. Methyl acrylate (15.9 g) was added to the solution, which was then warmed at about 40°C. Cuprous chloride (7.0 g) was added little by little to the mixture with vigorous stirring, and stirring was continued at the reaction temperature maintained at not higher than 45°C for about 3 hours until the evolution of nitrogen stopped. The

0155845

reaction solution was concentrated under reduced pressure, and the concentrate was made alkaline with aqueous concentrated ammonia and extracted with ethyl acetate. The ethyl acetate layer was washed with water, dried (MgSO_4) and freed of the solvent by distillation. The residue was chromatographed on a silica-gel (230 g) column, and elution was effected with ethyl ether-hexane-triethylamine (75:25:1, V/V) to give methyl 2-bromo-3-{4-[3-hydroxy-2-(3-methyl-2-pyridyl)propoxy]phenyl}-propionate in the form of an oily substance. Yield of 4.8 g.

IR (Neat): 3370, 1735 cm^{-1} . NMR (CDCl_3) δ : 2.35(3H,s), 2.9-3.4(2H,m), 3.3-3.7(1H,m), 3.63(3H,s), 3.9(1H,broad), 4.07(2H,d,J=3Hz), 4.1-4.6(3H,m), 6.76(2H,d,J=9Hz), 7.04(2H,d,J=9Hz), 7.3-7.6(2H,m), 8.30(1H,dd,J=5 and 2Hz).

Reference Example 3

- (1) By following the same procedure as described in Reference Example 1-(1), methyl 2-bromo-3-{4-[2-(5-ethyl-2-pyridyl)ethoxy]phenyl}propionate (18.5 g) was oxidized with m-chloroperbenzoic acid to give methyl 2-bromo-3-{4-[2-(5-ethyl-2-pyridyl)ethoxy]phenyl}propionate N-oxide. Yield of 19.2 g.
- (2) Methyl 2-bromo-3-{4-[2-(5-ethyl-2-pyridyl)ethoxy]phenyl}-propionate N-oxide (19.0 g) as obtained in Reference Example 3-(1) was reacted with acetic anhydride in the same manner as described in Reference Example 1-(2), and the resulting product was purified by silica-gel column chromatography to give methyl 3-{4-[2-acetoxy-2-(5-ethyl-2-pyridyl)ethoxy]phenyl}-2-bromopropionate in the form of an oily substance. Yield of 10.2 g. IR (Neat): 1730 cm^{-1} . NMR (CDCl_3) δ : 1.23(3H,t,J=7Hz), 2.13(3H,s), 2.66(2H,q,J=7Hz), 3.12(1H,dd,J=14 and 7Hz), 3.40(1H,dd,J=14 and 7Hz), 3.70(3H,s), 4.33(1H,t,J=7Hz), 4.40(2H,d,J=5Hz), 6.17(1H,t,J=5Hz), 6.83(2H,d,J=9Hz), 7.11(2H,d,J=9Hz), 7.33(1H,d,J=8Hz), 7.54(1H,dd,J=8 and 2Hz), 8.43(1H,d,J=2Hz).

Reference Example 4

- By following the same procedure as described in Reference Example 2, there were obtained the following compounds:

- (1) 4-[2-(5-Ethyl-2-pyridyl)-3-hydroxypropoxy]nitrobenzene:
an oily substance, IR (Neat): 3250, 1330 cm^{-1} . NMR (CDCl_3) δ :
1.23(3H,t,J=7Hz), 2.63(2H,q,J=7Hz), 3.35(1H,m), 4.10(2H,d,
J=4Hz), 4.50(3H,m), 6.93(2H,d,J=9Hz), 7.19(1H,d,J=8Hz),
5 7.51(1H,dd,J=8 and 2Hz), 8.13(2H,d,J=9Hz), 8.35(1H,d,J=2Hz).
(2) Methyl 2-bromo-3-{4-[2-(5-ethyl-2-pyridyl)-3-hydroxy-
propoxy]phenyl}propionate: an oily substance, IR (Neat):
3300, 1730 cm^{-1} . NMR (CDCl_3) δ : 1.23(3H,t,J=7Hz), 2.62(2H,q,
J=7Hz), 3.0 to 3.5(3H,m), 3.68(3H,s), 4.09(2H,d,J=4Hz), 4.2-
10 4.6(3H,m), 6.7-7.6(6H,m), 8.35(1H,d,J=2Hz).

Reference Example 5

By following the same procedure as described in
Reference Example 1, there were obtained the following
15 compounds:

- ° methyl 3-{4-[2-acetoxy-2-(2-pyridyl)ethoxy]phenyl}-2-
bromopropionate. Oily substance. Yield: 54.4%.

IR (Neat): 1740 cm^{-1}

NMR (CDCl_3) δ : 2.14(3H,s), 3.11(1H,dd,J=14 and 7Hz),
20 3.38(1H,dd,J=14 and 7Hz), 3.69(3H,s), 4.33(1H,t,
J=7Hz), 4.40(2H,d,J=5Hz), 6.17(1H,t,J=5Hz),
6.80(2H,d,J=9Hz), 8.08(2H,d,J=9Hz), 7.2~7.8(3H,m),
8.25(1H,dd,J=5 and 2Hz).

- ° methyl 3-{4-[2-acetoxy-2-(3-methyl-2-pyridyl)ethoxy]phenyl}-
25 2-bromopropionate. Oily substance. Yield: 30.1%

IR (Neat): 1730 cm^{-1}

NMR (CDCl_3) δ : 2.09(3H,s), 2.49(3H,s), 3.12(1H,dd,
J=14 and 7Hz), 3.40(1H,dd,J=14 and 7Hz), 3.70(3H,s),
4.1~4.6(3H,m), 6.33(1H,dd,J=8 and 5Hz), 6.81(2H,d,
J=9Hz), 8.08(2H,d,J=9Hz), 7.2~7.6(2H,m), 8.44(1H,
30 dd,J=5 and 2Hz).

- ° methyl 3-{4-[2-acetoxy-2-(5-methyl-2-pyridyl)ethoxy]phenyl}-
2-bromopropionate. Oily substance. Yield: 61.0%

IR (Neat): 1740 cm^{-1}

NMR (CDCl_3) δ : 2.19(3H,s), 2.30(3H,s), 3.10(1H,dd,
J=14 and 7Hz), 3.37(1H,dd,J=14 and 7Hz),
4.30(1H,t,J=7Hz), 4.37(2H,d,J=5Hz), 6.12(1H,t,J=5Hz),
6.79(2H,d,J=9Hz), 8.06(2H,d,J=9Hz), 7.2~7.6(2H,m),
8.37(2H,d,J=2Hz).

- ° methyl 3-{4-[2-acetoxy-2-(4,6-dimethyl-2-pyridyl)ethoxy]
phenyl}-2-bromopropionate. Oily substance. Yield: 51.2%

IR (Neat): 1740 cm^{-1}

NMR (CDCl_3) : 2.13(3H,s), 2.28(3H,s), 2.47(3H,s),
3.12(1H,dd,J=14 and 7Hz), 3.38(1H,dd,J=14 and 7Hz),
3.69(3H,s), 4.31(1H,t,J=7Hz), 4.37(2H,d,J=5Hz),
6.10(1H,t,J=5Hz), 6.7~7.2(6H,m).

Experiment Example

Blood sugar- and lipid-lowering activity in mice.


0.005 % of a test compound was mixed with a powder diet (CE-2, produced by CLEA JAPAN), and KKAY mice (male, 8 to 10 weeks of age, 5 mice/group) were placed on the diet for 4 days, and allowed free access to water throughout the experiment. Blood samples were taken from the orbital veins and assayed for blood sugar by the glucose oxidase method and for plasma triglyceride by the enzyme method in which the produced glycerol was determined by means of Cleantech TG-S Kit (Yatron), respectively. The respective values are expressed in terms of rate of lowering (%) from the values of control group not receiving the test compound.

Test compound (No. of Example)	Hypoglycemic activity (%)	Hypolipemic activity (%)
1	45	47
2	22	11
3	32	35
4	45	38
7	53	58
8	47	53
9	35	19
10	40	47

The novel thiazolidinedione derivatives (I) according to the present invention exhibit excellent blood sugar- and lipid-lowering activity, and are of value as pharmaceuticals, such as a therapeutic agent for diabetes and therapeutic agent for hyperlipemia, and so forth.


C L A I M S

1. A compound of the general formula:




[wherein R^1 and R^2 are the same or different and each represent hydrogen or a lower alkyl group; R^3 is hydrogen or an acyl group; n is 0 or 1] or a salt thereof.


2. A compound as claimed in Claim 1, wherein R^3 is hydrogen.
3. A compound as claimed in Claim 1, wherein n is 0.
4. A compound as claimed in Claim 1, wherein the compound is 5-[4-[2-hydroxy-2-(6-methyl-2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
5. A compound as claimed in Claim 1, wherein the compound is 5-[4-[2-(5-ethyl-2-pyridyl)-2-hydroxyethoxy]benzyl]-2,4-thiazolidinedione.
6. A compound as claimed in Claim 1, wherein the compound is 5-[4-[2-hydroxy-2-(3-methyl-2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
7. A compound as claimed in Claim 1, wherein the compound is 5-[4-[2-hydroxy-2-(2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
8. A process for producing a thiazolidinedione derivative of the general formula:



[wherein R^1 , R^2 and n are as defined hereinafter; R^3 is hydrogen or an acyl group], characterized in that said process comprises reacting a compound of the general formula:




[wherein R^1 and R^2 are the same or different and each represent hydrogen or a lower alkyl group; R^4 is hydrogen or an acyl group; R^5 is hydrogen or a lower alkyl group; X is a halogen atom; n is 0 or 1] with thiourea to produce a compound of the general formula:



[wherein each of the symbols is as defined hereinbefore] and hydrolyzing the resulting compound, followed by acylating if necessary.


9. A pharmaceutical composition which contains a thiazolidinedione derivative of the general formula:




[wherein R^1 and R^2 are the same or different and each represent hydrogen or a lower alkyl group; R^3 is hydrogen or an acyl group; n is 0 or 1] or a salt thereof.

AUSTRIAN CLAIMS


1. A process for producing a thiazolidinedione derivative of the general formula:



[wherein R^1 , R^2 and n are as defined hereinafter; R^3 is hydrogen or an acyl group], characterized in that said process comprises reacting a compound of the general formula:




[wherein R^1 and R^2 are the same or different and each represent hydrogen or a lower alkyl group; R^4 is hydrogen or an acyl group; R^5 is hydrogen or a lower alkyl group; X is a halogen atom; n is 0 or 1] with thiourea to produce a compound of the general formula:



[wherein each of the symbols is as defined hereinbefore] and hydrolyzing the resulting compound, followed by acylating if necessary.

2. A process for producing a thiazolidinedione derivative as claimed in Claim 1, wherein R^3 is hydrogen.
3. A process for producing a thiazolidinedione derivative as claimed in Claim 1, wherein n is 0.
4. A process for producing a thiazolidinedione derivative as claimed in Claim 1, wherein the thiazolidinedione derivative is 5-[4-[2-hydroxy-2-(6-methyl-2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
5. A process for producing a thiazolidinedione derivative as claimed in Claim 1, wherein the thiazolidinedione derivative is 5-[4-[2-(5-ethyl-2-pyridyl)-2-hydroxyethoxy]benzyl]-2,4-thiazolidinedione.
6. A process for producing a thiazolidinedione derivative as claimed in Claim 1, wherein the thiazolidinedione derivative is 5-[4-[2-hydroxy-2-(3-methyl-2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
7. A process for producing a thiazolidinedione derivative as claimed in Claim 1, wherein the thiazolidinedione derivative is 5-[4-[2-hydroxy-2-(2-pyridyl)ethoxy]benzyl]-2,4-thiazolidinedione.
8. A pharmaceutical composition which contains a thiazolidinedione derivative of the general formula:



[wherein R^1 and R^2 are the same or different and each represent hydrogen or a lower alkyl group; R^3 is hydrogen or an acyl group; n is 0 or 1] or a salt thereof.



European Patent
Office

EUROPEAN SEARCH REPORT

0155845

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 85301895.0
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	EP - A1 - 0 008 203 (TAKEDA) * Compound 40 *	1,8	C 07 D 417/12 A 61 K 31/44
A	CHIMICAL ABSTRACTS, vol. 98, no. 19, May 9, 1983, Columbus, Ohio, USA SOHDA, TAKASHI; MIZUNO, KATSUTOSHI; IMAMIYA, EIKO; TAWADA, HIROYUKI; MEGURO, KANJI; KAWAMATSU, YUTAKA; YAMAMOTO, YUJIRO "Studies on anti-diabetic agents. III. 5-Arylthiazolidine-2,4-diones as potent aldose reductase inhibitors" page 505, column 2, abstract-no. 160 624h & Chem. Pharm. Bull. 1982, 30(10), 3601-16	1	
A	CHIMICAL ABSTRACTS, vol. 98, no. 15, April 11, 1983, Columbus, Ohio, USA SOHDA, TAKASHI; MIZUNO, KATSUTOSHI; IMAMIYA, EIKO; SUGIYAMA, YASUO; FUJITA, TAKESHI; KAWAMATSU, YUTAKA "Studies on antidiabetic agents. II. Synthesis of 5-[4-(1-methylcyclohexylmethoxy)benzyl]thiazolidine-2,4-dione (ADD-3878) and its derivatives" page 627, column 2, abstract-no. 125 945s & Chem. Pharm. Bull. 1982, 30(10), 3580-600	1	TECHNICAL FIELDS SEARCHED (Int. Cl. 4) C 07 D 417/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 21-05-1985	Examiner HAMMER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



European Patent
Office

EUROPEAN SEARCH REPORT

0155845

Application number

- 2 -

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 85301895.0
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	<p>CHEMICAL ABSTRACTS, vol. 99, no. 23, December 5, 1983, Columbus, Ohio, USA</p> <p>SOHDA, TAKASHI; MIZUNO, KATSUTOSHI; HIRATA, TAKEO; MAKI, YOSHITAKA; KAWAMATSU, YUTAKA "Antiulcer activity of 5-benzylthiazolidine-2,4-dione derivatives" page 744, column 2, abstract-no. 194 854u</p> <p>& Chem. Pharm. Bull. 1983, 31(2), 560-9</p> <p>-----</p>	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 21-05-1985	Examiner HAMMER
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	